

A new estimate for Bochner-Riesz operators at the critical index on the weighted Hardy spaces

Hua Wang *

Department of Mathematics, Zhejiang University, Hangzhou 310027, China

Abstract

Let w be a Muckenhoupt weight and $H_w^p(\mathbb{R}^n)$ be the weighted Hardy spaces. In this paper, by using the atomic decomposition of $H_w^p(\mathbb{R}^n)$, we will show that the Bochner-Riesz operators T_R^δ are bounded from $H_w^p(\mathbb{R}^n)$ to the weighted weak Hardy spaces $WH_w^p(\mathbb{R}^n)$ when $0 < p < 1$ and $\delta = n/p - (n+1)/2$. This result is new even in the unweighted case.

MSC(2010): 42B15; 42B25; 42B30

Keywords: Bochner-Riesz operators; weighted Hardy spaces; weighted weak Hardy spaces; A_p weights; atomic decomposition

1 Introduction

The Bochner-Riesz operators of order $\delta > 0$ in $\mathbb{R}^n$ are defined initially for Schwartz functions in terms of Fourier transforms by

$$(\widehat{T_R^\delta f})(\xi) = \left(1 - \frac{|\xi|^2}{R^2}\right)_+^\delta \hat{f}(\xi), \quad 0 < R < \infty,$$

where $\hat{f}$ denotes the Fourier transform of f . The associated maximal Bochner-Riesz operator is defined by

$$T_*^\delta f(x) = \sup_{R>0} |T_R^\delta f(x)|.$$

These operators were first introduced by Bochner [1] in connection with summation of multiple Fourier series and played an important role in harmonic analysis. The problem concerning the spherical convergence of Fourier integrals have led to the study of their L^p boundedness. As for their H^p boundedness, Sjölin [17] and Stein, Taibleson and Weiss [18] proved the following theorem (see also [11, page 130]).

Theorem I. *Suppose that $0 < p \leq 1$ and $\delta > n/p - (n+1)/2$. Then there exists a constant $C > 0$ independent of f and R such that*

$$\|T_R^\delta(f)\|_{H^p} \leq C\|f\|_{H^p}.$$

*E-mail address: wanghua@pku.edu.cn.

In [18], the authors also considered weak type estimate for the maximal Bochner-Riesz operator T_*^δ at the critical index $\delta = n/p - (n+1)/2$ and showed the following inequality is sharp.

Theorem II. *Suppose that $0 < p < 1$ and $\delta = n/p - (n+1)/2$. Then there exists a constant $C > 0$ independent of f such that*

$$\sup_{\lambda > 0} \lambda^p |\{x \in \mathbb{R}^n : T_*^\delta f(x) > \lambda\}| \leq C \|f\|_{H^p}^p.$$

In 1995, Sato [15] studied the weighted case and obtained the following weighted weak type estimate for the maximal Bochner-Riesz operator T_*^δ .

Theorem III. *Let $w \in A_1$ (Muckenhoupt weight class), $0 < p < 1$ and $\delta = n/p - (n+1)/2$. Then there exists a constant $C > 0$ independent of f such that*

$$\sup_{\lambda > 0} \lambda^p \cdot w(\{x \in \mathbb{R}^n : T_*^\delta f(x) > \lambda\}) \leq C \|f\|_{H_w^p}^p.$$

In [16], Sato also showed that for $n \geq 2$, there exists a function $f \in H_w^1 \cap L^1$, $w \in A_1$, such that

$$\limsup_{R \rightarrow \infty} |T_R^{(n-1)/2} f(x)| = +\infty \quad \text{almost everywhere.}$$

Hence, $\delta = n/p - (n+1)/2$ is indeed the critical index for the weighted case.

In 2006, Lee [8] considered values of δ greater than the critical index $n/p - (n+1)/2$ and proved the $H_w^p - L_w^p$ boundedness of the maximal operator T_*^δ . Furthermore, by using this strong type estimate of T_*^δ , Lee [8] also obtained the H_w^p boundedness of the Bochner-Riesz operator.

The purpose of this article is to discuss the boundedness of Bochner-Riesz operators at the critical index $n/p - (n+1)/2$ on the weighted Hardy spaces. Our main result is stated as follows.

Theorem 1.1. *Let $0 < p < 1$, $\delta = n/p - (n+1)/2$ and $w \in A_1$. Suppose that $n(1/p - 1)$ is not a positive integer, then there exists a constant $C > 0$ independent of f and R such that*

$$\|T_R^\delta(f)\|_{WH_w^p} \leq C \|f\|_{H_w^p},$$

where WH_w^p denotes the weighted weak Hardy space.

In particular, if we take w to be a constant function, then we can get

Corollary 1.2. *Let $0 < p < 1$ and $\delta = n/p - (n+1)/2$. Suppose that $n(1/p - 1)$ is not a positive integer, then there exists a constant $C > 0$ independent of f and R such that*

$$\|T_R^\delta(f)\|_{WH^p} \leq C \|f\|_{H^p},$$

where WH^p denotes the weak Hardy space.

2 Notations and definitions

The definition of A_p class was first used by Muckenhoupt [13], Hunt, Muckenhoupt and Wheeden [7], and Coifman and Fefferman [2] in the study of weighted L^p boundedness of Hardy-Littlewood maximal functions and singular integrals. Let w be a nonnegative, locally integrable function defined on $\mathbb{R}^n$; all cubes are assumed to have their sides parallel to the coordinate axes. We say that $w \in A_p$, $1 < p < \infty$, if

$$\left(\frac{1}{|Q|} \int_Q w(x) dx \right) \left(\frac{1}{|Q|} \int_Q w(x)^{-1/(p-1)} dx \right)^{p-1} \leq C \quad \text{for every cube } Q \subseteq \mathbb{R}^n,$$

where C is a positive constant which is independent of the choice of Q .

For the case $p = 1$, $w \in A_1$, if

$$\frac{1}{|Q|} \int_Q w(x) dx \leq C \cdot \operatorname{ess\,inf}_{x \in Q} w(x) \quad \text{for every cube } Q \subseteq \mathbb{R}^n.$$

A weight function $w \in A_\infty$ if it satisfies the A_p condition for some $1 < p < \infty$. It is well known that if $w \in A_p$ with $1 < p < \infty$, then $w \in A_r$ for all $r > p$, and $w \in A_q$ for some $1 < q < p$. We thus write $q_w \equiv \inf\{q > 1 : w \in A_q\}$ to denote the critical index of w .

Given a cube Q and $\lambda > 0$, λQ denotes the cube with the same center as Q whose side length is λ times that of Q . $Q = Q(x_0, r)$ denotes the cube centered at x_0 with side length r . For a weight function w and a measurable set E , we denote the Lebesgue measure of E by $|E|$ and set the weighted measure $w(E) = \int_E w(x) dx$.

We give the following result that will be used in the sequel.

Lemma 2.1 ([6]). *Let $w \in A_1$. Then, for any cube Q , there exists an absolute constant $C > 0$ such that*

$$w(2Q) \leq C w(Q).$$

In general, for any $\lambda > 1$, we have

$$w(\lambda Q) \leq C \cdot \lambda^n w(Q),$$

where C does not depend on Q nor on λ .

Lemma 2.2 ([6]). *Let $w \in A_1$. Then there exists a constant $C > 0$ such that*

$$C \cdot \frac{|E|}{|Q|} \leq \frac{w(E)}{w(Q)}$$

for any measurable subset E of a cube Q .

Given a weight function w on $\mathbb{R}^n$, for $0 < p < \infty$, we denote by $L_w^p(\mathbb{R}^n)$ the weighted space of all functions satisfying

$$\|f\|_{L_w^p} = \left(\int_{\mathbb{R}^n} |f(x)|^p w(x) dx \right)^{1/p} < \infty.$$

We also denote by $WL_w^p(\mathbb{R}^n)$ the weighted weak L^p space which is formed by all functions satisfying

$$\|f\|_{WL_w^p} = \sup_{\lambda>0} \lambda \cdot w(\{x \in \mathbb{R}^n : |f(x)| > \lambda\})^{1/p} < \infty.$$

We write $\mathcal{S}(\mathbb{R}^n)$ to denote the Schwartz space of all rapidly decreasing infinitely differentiable functions and $\mathcal{S}'(\mathbb{R}^n)$ to denote the space of all tempered distributions, i.e., the topological dual of $\mathcal{S}(\mathbb{R}^n)$. For any $0 < p \leq 1$, the weighted Hardy spaces $H_w^p(\mathbb{R}^n)$ can be defined in terms of maximal functions. Let φ be a function in $\mathcal{S}(\mathbb{R}^n)$ satisfying $\int_{\mathbb{R}^n} \varphi(x) dx = 1$. Set

$$\varphi_t(x) = t^{-n} \varphi(x/t), \quad t > 0, x \in \mathbb{R}^n.$$

We will define the maximal function $M_\varphi^+ f(x)$ by

$$M_\varphi^+ f(x) = \sup_{t>0} |(\varphi_t * f)(x)|.$$

Then $H_w^p(\mathbb{R}^n)$ consists of those tempered distributions $f \in \mathcal{S}'(\mathbb{R}^n)$ for which $M_\varphi^+ f \in L_w^p(\mathbb{R}^n)$ with $\|f\|_{H_w^p} = \|M_\varphi^+ f\|_{L_w^p}$. The real-variable theory of weighted Hardy spaces has been investigated by many authors. For example, Garcia-Cuerva [5] studied the atomic decomposition and the dual spaces of H_w^p for $0 < p \leq 1$. The molecular characterization of H_w^p for $0 < p \leq 1$ was given by Lee and Lin [9]. We refer the readers to [5, 9, 20] and the references therein for further details.

Let us now turn to the weighted weak Hardy spaces, which are good substitutes for the weighted Hardy spaces in the study of the boundedness of some operators. The weak H^p spaces have first appeared in the work of Fefferman, Rivière and Sagher [3]. The atomic decomposition theory of weak H^1 space on $\mathbb{R}^n$ was given by Fefferman and Soria [4]. Later, Liu [10] established the weak H^p spaces on homogeneous groups. In 2000, Quek and Yang [14] introduced the weighted weak Hardy spaces and established their atomic decompositions. Let $w \in A_\infty$, $0 < p \leq 1$ and $N = [n(q_w/p - 1)]$. Define

$$\mathcal{A}_{N,w} = \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n) : \sup_{x \in \mathbb{R}^n} \sup_{|\alpha| \leq N+1} (1 + |x|)^{N+n+1} |D^\alpha \varphi(x)| \leq 1 \right\},$$

where $\alpha = (\alpha_1, \dots, \alpha_n) \in (\mathbb{N} \cup \{0\})^n$, $|\alpha| = \alpha_1 + \dots + \alpha_n$, and

$$D^\alpha \varphi = \frac{\partial^{|\alpha|} \varphi}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}}.$$

For $f \in \mathcal{S}'(\mathbb{R}^n)$, the nontangential grand maximal function of f is defined by

$$G_w f(x) = \sup_{\varphi \in \mathcal{A}_{N,w}} \sup_{|y-x|<t} |(\varphi_t * f)(y)|.$$

Then we can define the weighted weak Hardy space $WH_w^p(\mathbb{R}^n)$ by $WH_w^p(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G_w f \in WL_w^p(\mathbb{R}^n)\}$ and $\|f\|_{WH_w^p} = \|G_w f\|_{WL_w^p}$. In order to simplify the computations, we also use another equivalent definition of $WH_w^p(\mathbb{R}^n)$.

$$WH_w^p(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G_w^+ f \in WL_w^p(\mathbb{R}^n)\},$$

where $G_w^+ f$ is called the radial grand maximal function, which is defined by

$$G_w^+ f(x) = \sup_{\varphi \in \mathcal{A}_{N,w}} \sup_{t>0} |(\varphi_t * f)(x)|.$$

Moreover, we set $\|f\|_{WH_w^p} = \|G_w^+ f\|_{WL_w^p}$.

For w equals to a constant function, we shall denote $WL_w^p(\mathbb{R}^n)$, $H_w^p(\mathbb{R}^n)$ and $WH_w^p(\mathbb{R}^n)$ simply by $WL^p(\mathbb{R}^n)$, $H^p(\mathbb{R}^n)$ and $WH^p(\mathbb{R}^n)$.

Throughout this article C denotes a positive constant, which is independent of the main parameters and not necessarily the same at each occurrence.

3 The atomic decomposition for weighted Hardy spaces

In this article, we will use Garcia-Cuerva's atomic decomposition theory for weighted Hardy spaces in [5, 20]. We characterize weighted Hardy spaces in terms of atoms in the following way.

Let $0 < p \leq 1 \leq q \leq \infty$ and $p \neq q$ such that $w \in A_q$ with critical index q_w . Set $[\cdot]$ the greatest integer function. For $s \in \mathbb{Z}_+$ satisfying $s \geq N = [n(q_w/p - 1)]$, a real-valued function $a(x)$ is called a (p, q, s) -atom centered at x_0 with respect to w (or a w -(p, q, s)-atom centered at x_0) if the following conditions are satisfied:

- (a) $a \in L_w^q(\mathbb{R}^n)$ and is supported in a cube Q centered at x_0 ;
- (b) $\|a\|_{L_w^q} \leq w(Q)^{1/q-1/p}$;
- (c) $\int_{\mathbb{R}^n} a(x) x^\alpha dx = 0$ for every multi-index α with $|\alpha| \leq s$.

Theorem 3.1. *Let $0 < p \leq 1 \leq q \leq \infty$ and $p \neq q$ such that $w \in A_q$ with critical index q_w . For each $f \in H_w^p(\mathbb{R}^n)$, there exist a sequence $\{a_j\}$ of w -(p, q, N)-atoms and a sequence $\{\lambda_j\}$ of real numbers with $\sum_j |\lambda_j|^p \leq C \|f\|_{H_w^p}^p$ such that $f = \sum_j \lambda_j a_j$ both in the sense of distributions and in the H_w^p norm.*

4 Proof of Theorem 1.1

The Bochner-Riesz operators can be expressed as convolution operators

$$T_R^\delta f(x) = (\phi_{1/R} * f)(x),$$

where $\phi(x) = [(1 - |\cdot|^2)_+]^\delta(x)$ and $\phi_{1/R}(x) = R^n \phi(Rx)$. It is well known that the kernel ϕ can be represented as (see [12, 19])

$$\phi(x) = \pi^{-\delta} \Gamma(\delta + 1) |x|^{-(\frac{n}{2} + \delta)} J_{\frac{n}{2} + \delta}(2\pi|x|),$$

where $J_\mu(t)$ is the Bessel function

$$J_\mu(t) = \frac{(\frac{t}{2})^\mu}{\Gamma(\mu + \frac{1}{2})\Gamma(\frac{1}{2})} \int_{-1}^1 e^{its} (1 - s^2)^{\mu - \frac{1}{2}} ds.$$

The following kernel estimates of these convolution operators are well known. For its proof, we refer the readers to [15]. See also [11, page 121].

Lemma 4.1. *Let $0 < p < 1$ and $\delta = n/p - (n+1)/2$. Then the kernel ϕ satisfies the inequality*

$$\sup_{x \in \mathbb{R}^n} (1 + |x|)^{n/p} |D^\alpha \phi(x)| \leq C \quad \text{for all multi-indices } \alpha.$$

In order to prove our main theorem, we shall need the following superposition principle on weighted weak type estimates.

Lemma 4.2. *Let $w \in A_1$ and $0 < p < 1$. If a sequence of measurable functions $\{f_j\}$ satisfy*

$$w(\{x \in \mathbb{R}^n : |f_j(x)| > \alpha\}) \leq \alpha^{-p} \quad \text{for all } j \in \mathbb{Z}$$

and

$$\sum_{j \in \mathbb{Z}} |\lambda_j|^p \leq 1,$$

then we have that $\sum_j \lambda_j f_j(x)$ is absolutely convergent almost everywhere and

$$w\left(\left\{x \in \mathbb{R}^n : \left|\sum_j \lambda_j f_j(x)\right| > \alpha\right\}\right) \leq \frac{2-p}{1-p} \cdot \alpha^{-p}.$$

Proof. The proof of this lemma is similar to the corresponding result for the unweighted case which can be found in [18]. See also [11, page 123]. $\square$

We now prove the following estimate which plays a key role in the proof of our main result.

Lemma 4.3. *Let $0 < p < 1 < q < \infty$, $\delta = n/p - (n+1)/2$ and $w \in A_1$. Then for any given w -(p, q, N)-atom a with support contained in $Q = Q(x_0, r)$, we have*

$$T_*^\delta(a)(y) \leq C \cdot \frac{r^{n/p}}{|y - x_0|^{n/p}} w(Q)^{-1/p} \quad \text{whenever } |y - x_0| > \sqrt{n} r.$$

Here, and in what follows, we always denote $N = [n(q_w/p - 1)] = [n(1/p - 1)]$.

Proof. Actually, this lemma was essentially contained in [8]. For the sake of completeness, we give its proof here. For any $\varepsilon > 0$, we write

$$(\phi_\varepsilon * a)(y) = \varepsilon^{-n} \int_{Q(x_0, r)} \phi\left(\frac{y - z}{\varepsilon}\right) a(z) dz.$$

Let us now consider the following two cases.

(i) $0 < \varepsilon \leq r$. Note that $\delta = n/p - (n+1)/2$, then by Lemma 4.1, we have

$$|(\phi_\varepsilon * a)(y)| \leq C \cdot \varepsilon^{n/p - n} \int_{Q(x_0, r)} \frac{|a(z)|}{|y - z|^{n/p}} dz.$$

By our assumption, when $z \in Q(x_0, r)$, then we can easily get $|y - z| \geq |y - x_0| - |z - x_0| \geq \frac{|y - x_0|}{2}$. Observe that $0 < p < 1$, then $n/p - n > 0$. Hence

$$|(\phi_\varepsilon * a)(y)| \leq C \cdot \frac{r^{n/p-n}}{|y - x_0|^{n/p}} \int_{Q(x_0, r)} |a(z)| dz. \quad (1)$$

Denote the conjugate exponent of $q > 1$ by $q' = q/(q - 1)$. Using Hölder's inequality, the A_q condition and the size condition of atom a , we can get

$$\begin{aligned} \int_{Q(x_0, r)} |a(z)| dz &\leq \left(\int_{Q(x_0, r)} |a(z)|^q w(z) dz \right)^{1/q} \left(\int_{Q(x_0, r)} (w(z)^{-1/q})^{q'} dz \right)^{1/q'} \\ &\leq C \cdot \|a\|_{L_w^q} |Q(x_0, r)| w(Q)^{-1/q} \\ &\leq C \cdot |Q(x_0, r)| w(Q)^{-1/p}. \end{aligned} \quad (2)$$

Substituting the above inequality (2) into (1), we thus obtain

$$|(\phi_\varepsilon * a)(y)| \leq C \cdot \frac{r^{n/p}}{|y - x_0|^{n/p}} w(Q)^{-1/p}. \quad (3)$$

(ii) $\varepsilon > r$. It is easy to see that the choice of N in this lemma implies $\frac{n}{n+N+1} < p \leq \frac{n}{n+N}$. Using the vanishing moment condition of atom a , Taylor's theorem and Lemma 4.1, we deduce

$$\begin{aligned} |(\phi_\varepsilon * a)(y)| &= \varepsilon^{-n} \left| \int_{Q(x_0, r)} \left[\phi\left(\frac{y-z}{\varepsilon}\right) - \sum_{|\gamma| \leq N} \frac{D^\gamma \phi\left(\frac{y-x_0}{\varepsilon}\right)}{\gamma!} \left(\frac{z-x_0}{\varepsilon}\right)^\gamma \right] a(z) dz \right| \\ &\leq \varepsilon^{-n} \cdot \left(\frac{\sqrt{n}r}{2\varepsilon}\right)^{N+1} \int_{Q(x_0, r)} \sum_{|\gamma|=N+1} \left| \frac{D^\gamma \phi\left(\frac{y-x_0-\theta(z-x_0)}{\varepsilon}\right)}{\gamma!} \right| |a(z)| dz \\ &\leq C \cdot \frac{r^{N+1}}{\varepsilon^{n+N+1}} \int_{Q(x_0, r)} \left| \frac{y-x_0-\theta(z-x_0)}{\varepsilon} \right|^{-n/p} |a(z)| dz, \end{aligned}$$

where $0 < \theta < 1$. As in the first case (i), we have $|y - x_0| \geq 2|z - x_0|$, which implies $|y - x_0 - \theta(z - x_0)| \geq |y - x_0| - |z - x_0| \geq \frac{|y - x_0|}{2}$. This together with the inequality (2) yields

$$|(\phi_\varepsilon * a)(y)| \leq C \cdot \frac{r^{n+N+1}}{\varepsilon^{n+N+1-n/p}} \frac{1}{|y - x_0|^{n/p}} w(Q)^{-1/p}.$$

Observe that $n + N + 1 - n/p > 0$, then for $\varepsilon > r$, we have $\varepsilon^{n+N+1-n/p} > r^{n+N+1-n/p}$. Consequently

$$|(\phi_\varepsilon * a)(y)| \leq C \cdot \frac{r^{n/p}}{|y - x_0|^{n/p}} w(Q)^{-1/p}. \quad (4)$$

Summarizing the estimates (3) and (4) derived above and taking the supremum over all $\varepsilon > 0$, we obtain the desired inequality. $\square$

We are now in a position to give the proof of Theorem 1.1.

Proof. By the atomic decomposition theorem (Theorem 3.1) and Lemma 4.2, it suffices to show that for any w -(p, q, N)-atom a , there exists a constant $C > 0$ independent of a such that $\|G_w^+(T_R^\delta a)\|_{WL_w^p} \leq C$.

Let a be a w -(p, q, N)-atom with $\text{supp } a \subseteq Q = Q(x_0, r)$, $0 < p < 1 < q < \infty$, and let $Q^* = (4\sqrt{n})Q$. For any given $\lambda > 0$, we write

$$\begin{aligned} & \lambda^p \cdot w(\{x \in \mathbb{R}^n : |G_w^+(T_R^\delta a)(x)| > \lambda\}) \\ & \leq \lambda^p \cdot w(\{x \in Q^* : |G_w^+(T_R^\delta a)(x)| > \lambda\}) + \lambda^p \cdot w(\{x \in (Q^*)^c : |G_w^+(T_R^\delta a)(x)| > \lambda\}) \\ & = I_1 + I_2. \end{aligned}$$

Let us deal with the term I_1 first. Applying Chebyshev's inequality and Hölder's inequality, we obtain

$$\begin{aligned} I_1 & \leq \int_{Q^*} |G_w^+(T_R^\delta a)(x)|^p w(x) dx \\ & \leq \left(\int_{Q^*} |G_w^+(T_R^\delta a)(x)|^q w(x) dx \right)^{p/q} \left(\int_{Q^*} w(x) dx \right)^{1-p/q} \\ & \leq \|G_w^+(T_R^\delta a)\|_{L_w^q}^p w(Q^*)^{1-p/q}. \end{aligned}$$

Since $0 < p < 1$ and $\delta = n/p - (n+1)/2$, then $\delta > (n-1)/2$. In this case, it is well known that (see [12, 19])

$$T_*^\delta a(x) \leq C \cdot Ma(x),$$

where M denotes the Hardy-Littlewood maximal operator. In addition, we can easily see that for any function f , the following inequality holds

$$G_w^+ f(x) \leq C \cdot Mf(x).$$

Since $w \in A_1$, then $w \in A_q$ for any $1 < q < \infty$. Hence, it follows from the L_w^q boundedness of M and Lemma 2.1 that

$$\begin{aligned} I_1 & \leq C \cdot \|T_*^\delta(a)\|_{L_w^q}^p w(Q^*)^{1-p/q} \\ & \leq C \cdot \|a\|_{L_w^q}^p w(Q)^{1-p/q} \\ & \leq C. \end{aligned} \tag{5}$$

We now turn to estimate the other term I_2 . First, we claim that for every multi-index γ with $|\gamma| \leq N$ and for any $0 < R < \infty$, the following equality holds

$$\int_{\mathbb{R}^n} T_R^\delta a(y) y^\gamma dy = 0. \tag{6}$$

In fact, by using the Leibnitz formula and vanishing moment condition of atom a , we thus have

$$\begin{aligned}
\int_{\mathbb{R}^n} T_R^\delta a(y) y^\gamma dy &= (T_R^\delta a(y) y^\gamma)^\wedge(0) \\
&= C \cdot D^\gamma(\widehat{T_R^\delta a})(0) \\
&= C \cdot D^\gamma(\widehat{\phi_{1/R} \cdot \widehat{a}})(0) \\
&= C \cdot \sum_{|\alpha|+|\beta|=|\gamma|} (D^\alpha \widehat{\phi_{1/R}})(0) \cdot (D^\beta \widehat{a})(0) \\
&= 0,
\end{aligned}$$

which is the desired conclusion. Then by the equality (6) derived above, we can split the following expression into three parts and estimate each term respectively.

$$\begin{aligned}
|\varphi_t * (T_R^\delta a)(x)| &= \left| \int_{\mathbb{R}^n} t^{-n} \left[\varphi\left(\frac{x-y}{t}\right) - \sum_{|\gamma| \leq N} \frac{D^\gamma \varphi\left(\frac{x-x_0}{t}\right) \left(\frac{y-x_0}{t}\right)^\gamma}{\gamma!} \right] T_R^\delta a(y) dy \right| \\
&\leq t^{-n} \left| \int_{|y-x_0| \leq \sqrt{n}r} \left[\varphi\left(\frac{x-y}{t}\right) - \sum_{|\gamma| \leq N} \frac{D^\gamma \varphi\left(\frac{x-x_0}{t}\right) \left(\frac{y-x_0}{t}\right)^\gamma}{\gamma!} \right] T_R^\delta a(y) dy \right| \\
&\quad + t^{-n} \left| \int_{\sqrt{n}r < |y-x_0| \leq \frac{|x-x_0|}{2}} \left[\varphi\left(\frac{x-y}{t}\right) - \sum_{|\gamma| \leq N} \frac{D^\gamma \varphi\left(\frac{x-x_0}{t}\right) \left(\frac{y-x_0}{t}\right)^\gamma}{\gamma!} \right] T_R^\delta a(y) dy \right| \\
&\quad + t^{-n} \left| \int_{|y-x_0| > \frac{|x-x_0|}{2}} \left[\varphi\left(\frac{x-y}{t}\right) - \sum_{|\gamma| \leq N} \frac{D^\gamma \varphi\left(\frac{x-x_0}{t}\right) \left(\frac{y-x_0}{t}\right)^\gamma}{\gamma!} \right] T_R^\delta a(y) dy \right| \\
&= J_1 + J_2 + J_3.
\end{aligned}$$

For the term J_1 , by using Taylor's theorem, we then obtain

$$J_1 \leq t^{-n} \cdot \left(\frac{\sqrt{n}r}{t}\right)^{N+1} \int_{|y-x_0| \leq \sqrt{n}r} \sum_{|\gamma|=N+1} \left| \frac{D^\gamma \varphi\left(\frac{x-x_0-\theta'(y-x_0)}{t}\right)}{\gamma!} \right| |T_*^\delta a(y)| dy,$$

where $0 < \theta' < 1$. For any $x \in (Q^*)^c$ and $|y-x_0| \leq \sqrt{n}r$, as before, we have $|x-x_0-\theta'(y-x_0)| \geq \frac{|x-x_0|}{2}$. Observe that $\varphi \in \mathcal{A}_{N,w}$, then it follows from Hölder's inequality and the size condition of atom a that

$$\begin{aligned}
J_1 &\leq C \cdot \frac{r^{N+1}}{t^{n+N+1}} \int_{|y-x_0| \leq \sqrt{n}r} \left| \frac{x-x_0-\theta'(y-x_0)}{t} \right|^{-n-N-1} |T_*^\delta a(y)| dy \\
&\leq C \cdot \frac{r^{N+1}}{|x-x_0|^{n+N+1}} \|T_*^\delta(a)\|_{L_w^q} |Q(x_0, 2\sqrt{n}r)| w(Q(x_0, 2\sqrt{n}r))^{-1/q} \\
&\leq C \cdot \frac{r^{N+1}}{|x-x_0|^{n+N+1}} \|a\|_{L_w^q} |Q(x_0, r)| w(Q)^{-1/q} \\
&\leq C \cdot \frac{r^{n+N+1}}{|x-x_0|^{n+N+1}} w(Q)^{-1/p}.
\end{aligned}$$

On the other hand, for the term J_2 , notice that $|y - x_0| > \sqrt{nr}$ and $\varphi \in \mathcal{A}_{N,w}$, then by Taylor's theorem and Lemma 4.3, we deduce

$$\begin{aligned} J_2 &\leq t^{-n} \int_{\sqrt{nr} < |y-x_0| \leq \frac{|x-x_0|}{2}} \sum_{|\gamma|=N+1} \left| \frac{D^\gamma \varphi\left(\frac{x-x_0-\theta'(y-x_0)}{t}\right)}{\gamma!} \right| \left| \frac{y-x_0}{t} \right|^{N+1} |T_*^\delta a(y)| dy \\ &\leq C \cdot t^{-n} \int_{\sqrt{nr} < |y-x_0| \leq \frac{|x-x_0|}{2}} \frac{t^{n+N+1}}{|x-x_0|^{n+N+1}} \cdot \frac{|y-x_0|^{N+1}}{t^{N+1}} \cdot \frac{r^{n/p}}{|y-x_0|^{n/p}} w(Q)^{-1/p} dy \\ &\leq C \cdot \frac{r^{n/p}}{|x-x_0|^{n+N+1}} w(Q)^{-1/p} \int_{\sqrt{nr} < |y-x_0| \leq \frac{|x-x_0|}{2}} |y-x_0|^{N+1-n/p} dy. \end{aligned}$$

If we rewrite this integral in polar coordinates and note the fact that $-1 < n + N - n/p \leq 0$, then we can get

$$\begin{aligned} \int_{\sqrt{nr} < |y-x_0| \leq \frac{|x-x_0|}{2}} |y-x_0|^{N+1-n/p} dy &\leq C \cdot \int_{\sqrt{nr}}^{\frac{|x-x_0|}{2}} \rho^{n+N-n/p} d\rho \\ &\leq C \cdot |x-x_0|^{n+N+1-n/p}. \end{aligned}$$

So we have

$$J_2 \leq C \cdot \frac{r^{n/p}}{|x-x_0|^{n/p}} w(Q)^{-1/p}.$$

For the last term J_3 , when $x \in (Q^*)^c$, then it is easy to see that $|y - x_0| > \frac{|x-x_0|}{2} > \sqrt{nr}$. Therefore, it follows from Lemma 4.3 and the fact $\varphi \in \mathcal{A}_{N,w}$ that

$$\begin{aligned} J_3 &\leq t^{-n} \int_{|y-x_0| > \frac{|x-x_0|}{2}} \left\{ \left| \varphi\left(\frac{x-y}{t}\right) \right| + \sum_{j=0}^N \sum_{|\gamma|=j} \left| \frac{D^\gamma \varphi\left(\frac{x-x_0}{t}\right)}{\gamma!} \right| \left| \frac{y-x_0}{t} \right|^j \right\} |T_*^\delta a(y)| dy \\ &\leq C \int_{|y-x_0| > \frac{|x-x_0|}{2}} \left\{ |\varphi_t(x-y)| + \sum_{j=0}^N \frac{|y-x_0|^j}{|x-x_0|^{n+j}} \right\} \cdot \frac{r^{n/p}}{|y-x_0|^{n/p}} w(Q)^{-1/p} dy \\ &\leq C \cdot w(Q)^{-1/p} \left(\frac{r^{n/p}}{|x-x_0|^{n/p}} \|\varphi_t\|_{L^1} + \sum_{j=0}^N \frac{r^{n/p}}{|x-x_0|^{n+j}} \int_{|y-x_0| > \frac{|x-x_0|}{2}} \frac{dy}{|y-x_0|^{n/p-j}} \right). \end{aligned}$$

By our assumption (say $n(1/p - 1)$ is not a positive integer), we know that $n/p - n > N = [n(1/p - 1)]$. Thus we have $n/p - n - j > 0$ for any $0 \leq j \leq N$. Making use of the polar coordinates again, we obtain

$$\begin{aligned} \int_{|y-x_0| > \frac{|x-x_0|}{2}} \frac{dy}{|y-x_0|^{n/p-j}} &\leq C \cdot \int_{\frac{|x-x_0|}{2}}^{\infty} \frac{1}{\rho^{n/p-n-j+1}} d\rho \\ &\leq C \cdot \frac{1}{|x-x_0|^{n/p-n-j}}. \end{aligned}$$

Hence

$$J_3 \leq C \cdot \frac{r^{n/p}}{|x-x_0|^{n/p}} w(Q)^{-1/p}.$$

Therefore, combining the above estimates for J_1 , J_2 and J_3 , and then taking the supremum over all $t > 0$ and all $\varphi \in \mathcal{A}_{N,w}$, we finally obtain that for any $x \in (Q^*)^c$,

$$\begin{aligned} |G_w^+(T_R^\delta a)(x)| &\leq C \cdot \left(\frac{r^{n+N+1}}{|x-x_0|^{n+N+1}} w(Q)^{-1/p} + \frac{r^{n/p}}{|x-x_0|^{n/p}} w(Q)^{-1/p} \right) \\ &\leq C \cdot \frac{r^{n/p}}{|x-x_0|^{n/p}} w(Q)^{-1/p}. \end{aligned} \quad (7)$$

Without loss of generality, we may assume that the absolute constant C appearing in the above inequality is bigger than one. For any fixed $\lambda > 0$, we are going to consider two cases. If $\lambda \geq C \cdot \frac{r^{n/p}}{(2\sqrt{nr})^{n/p}} w(Q)^{-1/p}$, then for any $x \in (Q^*)^c$, we have $|x-x_0| \geq 2\sqrt{nr}$. Thus in view of (7), we can easily see that $\{x \in (Q^*)^c : |G_w^+(T_R^\delta a)(x)| > \lambda\} = \emptyset$. Hence, in this case, the inequality

$$I_2 \leq C$$

holds trivially. Now suppose that $\lambda < C \cdot \frac{r^{n/p}}{(2\sqrt{nr})^{n/p}} w(Q)^{-1/p}$. If we set

$$R = \left(\frac{C^p \cdot |Q|}{\lambda^p \cdot w(Q)} \right)^{1/n},$$

then it is not difficult to verify that $R \geq 2\sqrt{nr} \geq r$ and

$$\{x \in (Q^*)^c : |G_w^+(T_R^\delta a)(x)| > \lambda\} \subseteq \{x \in \mathbb{R}^n : |x-x_0| < R\} \subseteq Q(x_0, 2R).$$

Since $w \in A_1$, then by Lemma 2.2, we can get (below, $\tilde{C}$ is an absolute constant)

$$\tilde{C} \cdot \frac{|Q(x_0, r)|}{|Q(x_0, R)|} \leq \frac{w(Q(x_0, r))}{w(Q(x_0, R))},$$

which is equivalent to

$$\begin{aligned} w(Q(x_0, R)) &\leq \frac{R^n \cdot w(Q)}{\tilde{C} \cdot |Q|} \\ &\leq \frac{C^p}{\tilde{C} \cdot \lambda^p}. \end{aligned}$$

Therefore

$$\begin{aligned} I_2 &\leq \lambda^p \cdot w(Q(x_0, 2R)) \\ &\leq C^p \cdot \tilde{C}^{-1}. \end{aligned} \quad (8)$$

Combining the above estimate (8) with (5) and then taking the supremum over all $\lambda > 0$, we complete the proof of Theorem 1.1. $\square$

Remark. In the end, we would like to point out that the conclusions of Theorem 1.1 and Corollary 1.2 are not generally true for the maximal Bochner-Riesz operators T_*^δ , because we can not deduce that T_*^δ satisfy the previous vanishing moment condition (6).

References

- [1] S. Bochner, Summation of multiple Fourier series by spherical means, Trans. Amer. Math. Soc, **40**(1936), 175–207.
- [2] R. R. Coifman and C. Fefferman, Weighted norm inequalities for maximal functions and singular integrals, Studia Math, **51**(1974), 241–250.
- [3] C. Fefferman, N. Rivière and Y. Sagher, Interpolation between H^p spaces: The real method, Trans. Amer. Math. Soc, **191**(1974), 75–81.
- [4] R. Fefferman and F. Soria, The space weak H^1 , Studia Math, **85**(1987), 1–16.
- [5] J. Garcia-Cuerva, Weighted H^p spaces, Dissertations Math, **162**(1979), 1–63.
- [6] J. Garcia-Cuerva and J. Rubio de Francia, Weighted Norm Inequalities and Related Topics, North-Holland, Amsterdam, 1985.
- [7] R. A. Hunt, B. Muckenhoupt and R. L. Wheeden, Weighted norm inequalities for the conjugate function and Hilbert transform, Trans. Amer. Math. Soc, **176**(1973), 227–251.
- [8] M. Y. Lee, Weighted norm inequalities of Bochner-Riesz means, J. Math. Anal. Appl, **324**(2006), 1274–1281.
- [9] M. Y. Lee and C. C. Lin, The molecular characterization of weighted Hardy spaces, J. Func. Anal, **188**(2002), 442–460.
- [10] H. P. Liu, The weak H^p spaces on homogeneous groups, Lecture Notes in Math, Vol. 1494, Springer-Verlag, 1991, 113–118.
- [11] S. Z. Lu, Four Lectures on Real H^p Spaces, World Scientific Publishing, River Edge, N.J., 1995.
- [12] S. Z. Lu and K. Y. Wang, Bochner-Riesz means(in Chinese), Beijing Normal Univ Press, Beijing, 1988.
- [13] B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, Trans. Amer. Math. Soc, **165**(1972), 207–226.
- [14] T. S. Quek and D. C. Yang, Calderón-Zygmund-type operators on weighted weak Hardy spaces over $\mathbb{R}^n$, Acta Math. Sinica, **16**(2000), 141–160.
- [15] S. Sato, Weak type estimates for some maximal operators on the weighted Hardy spaces, Ark. Mat., **33**(1995), 377–384.
- [16] S. Sato, Divergence of the Bochner-Riesz means in the weighted Hardy spaces, Studia Math, **118**(1996), 261–275.

- [17] P. Sjölin, Convolution with oscillating kernels in H^p spaces, J. London Math. Soc, **23**(1981), 442–454.
- [18] E. M. Stein, M. H. Taibleson and G. Weiss, Weak type estimates for maximal operators on certain H^p classes, Rend. Circ. Mat. Palermo, Suppl. 1, **2**(1981), 81–97.
- [19] E. M. Stein and G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, Princeton Univ. Press, Princeton, New Jersey, 1971.
- [20] J. O. Stömberg and A. Torchinsky, Weighted Hardy spaces, Lecture Notes in Math, Vol 1381, Springer-Verlag, 1989.